

## HerbVar - Reproductive Damage Protocol

By Phil Hahn (hahnp@ufl.edu), Susan Whitehead (swhitehead@vt.edu), Paul Ode (paul.ode@colostate.edu), and Will Wetzel (wcwetzel@msu.edu)

\*Updated 08 November 2019

*Please send any comments or suggestions you have regarding the protocol to the above people.*

**Overview:** This protocol aims to assess damage by insect herbivores to reproductive parts of plants (i.e., flowers, fruits, and/or seeds). This is a supplement to the [Primary HerbVar Survey Protocol](#), which aims to randomly select and sample 30 plants, plus their nearest conspecific neighbor, within a population.

**Objectives:** The goal is to measure the proportional damage to reproductive organs on each plant within the surveyed population. That is, for each individual plant we will record the number of damaged and undamaged reproductive organs. Ideally, these measurements should be taken as supplemental data for the same individual plants (focal and neighbor plants) for which leaf damage was taken for the primary protocol, although reproductive damage measurements are welcome from other plant populations from which no leaf damage measurements have been taken.

**What species to survey:** We are hoping to get broad taxonomic and geographic coverage of damage to reproductive organs. Therefore, any species could be surveyed. However, to ensure that the data are comparable across sites/species/families/etc, the plants should have the following characteristics:

- At least half of the individuals at your site should possess reproductive material. If most of the plants are in a vegetative stage, you probably won't be able to survey enough reproductive individuals to get a decent sample size. Ideally, >30 of the sampled plants will have reproductive damage data.
- Each individual plant should produce enough flowers/fruits/seeds so that you can survey between 15-30 reproductive units per plant. These 'units' could be flowers, fruits, or seeds, whichever will give you enough things to count. E.g., if your plants have just a few fruits, try opening fruits and counting a random sample of seeds, which could get you higher numbers. If some plants have fewer units (~5-10 flowers/fruits/seeds), that is okay, as long as most have at least 15.
- Most of the plants should be in a similar phenological stage. If there is a mixture of flowering and fruiting plants within the population, it might be difficult to get a large enough sample size for one organ type. Additionally, different phenological stages will likely be attacked by different insects.

If your plant does not meet these requirements, please skip measuring damage to reproductive organs. That's okay. Or get in touch if you have questions.

**Damage Measurements:** Overall we are looking to count the number of flowers/fruits/seeds with: 1) herbivore damage, 2) pathogen damage, 3) unknown damage, and 4) total number of units examined (see Table 1). Ideally at least 30 reproductive units will be examined per plant. For plants that have much more than 30 units, examine a haphazard subsample of 30 units per plant. These should be sampled from different parts of the plant to obtain a good representative sample of damage levels to the plant. Damage measurements will be highly variable among species, depending on the types of flowers or fruits produced as well as the type of damage that is most common in the population. Below we give some general guidelines. There likely will need to be modifications for some species, but we trust collaborators to do the best they can in their systems while maintaining the overall spirit of the protocol. Keep in mind that the goal is to capture the variability in damage rates among plants within a population, so you will want to choose a measure of flower/fruit/seed damage that best captures this variability.

**What to look for?** Start by surveying the type of damage that is most common on the reproductive structures of plants in the population. Look for damage by insects that may chew on developing flowers (e.g. katydids, beetles), insects that bore into flower heads or seeds (e.g. larval weevils, leps, or flies), or true bugs that may probe/pierce into the seeds or fruits (looks like little black dots on the fruits). In many cases you will need to tear open the seed head/fruit to look for boring insects inside the seeds/fruit. Look for signs of chewing damage inside the fruit, such as destroyed seeds and insect frass. The best measurements of damage will depend on the type and extent of damage present. Record the following data by adding some or all of these potential columns:

- #repro\_chew\_dmg (number of reproductive organs with chewing damage):
- #repro\_pierce\_dmg (number of reproductive organs with piercing damage):
- #repro\_path\_dmg (number of reproductive organs with pathogen damage):
- #repro\_unk\_dmg (number of reproductive organs with unknown damage):
- #repro\_total (total number of reproductive organs examined for damage):
- If present and identifiable, record the number and identity of the florivore, frugivore, seed predator

Table 1. Example of columns to add to the main data sheet.

Plant	#repro_chew_dmg	#repro_pierce_dmg	#repro_path_dmg	#repro_unk_dmg	#repro_total	Phenology
1	10	0	8	2	19	Fruiting
2	5	8	10	0	26	Fruiting
3	2	0	0	0	21	Fruiting
4						Vegetative

Note: #repro\_total can be greater than sum of others because units can have both herbivore and pathogen damage.

Examples of different types of damage to reproductive material



*Abronia umbellata* with chewing damage to two corolla (one is completely chewed and one is partially chewed). In this example, each floret would be scored for damage (1 damaged, 13 organs examined). Photo: Eric LoPresti.



*Vicia americana* seed pod. Notice boring holes in the upper-left seed and frass in the pod. In this example, each of 30 seeds would be scored for damage. Photo: Phil Hahn.



*Abronia turbinata* with damage and frass from a moth caterpillar (*Neogrotella macdunnoughi*). Note chewing damage to petals/corollas and caterpillars at base of florets. In this example, each of 30 florets would be scored for damage. Photo: Eric LoPresti.



*Monarda fistulosa* seed head ripped open to reveal a weevil larva. Notice the damage holes to the floral tubes. In this example, each flower head would be recorded as damaged (or undamaged) and ideally 30 flower heads would be assessed per plant. Often this species does not produce 30 flowers per plant, so smaller numbers would be acceptable. If most plants have fewer than 10 flowers, this would not be a good population to survey for reproductive damage. Photo: Phil Hahn.



*Lonicera* fruit with chewing damage. In this example, each of 30 fruits would be scored for damage. Photo: Susan Whitehead.



*Lonicera* fruit with piercing damage. In this example, each of 30 fruits would be scored for damage. Photo: Susan Whitehead.



*Aquilegia shockleyi* with chewing damage to fruits (from *Heliothis phloxiphaga*). In this example, each fruit could be scored for damage, or each fruit could be opened and seeds could be counted (infer missing, fully-consumed seeds from pod features). Photo: Eric LoPresti.